

Familial Mediterranean fever: What associations to screen for?

Salem Bouomrani, Ines Masmoudi, Sawssan Ben Teber

Sfax Faculty of Medicine, University of Sfax, Tunisia

Department of Internal Medicine, Military Hospital of Gabes, Tunisia

Abstract

Familial Mediterranean fever (FMF) is the most common and best known of hereditary recurrent fever or periodic fever syndromes. It was described in 1945 and genetically characterized in 1992. It is caused by a point mutation in the *MEFV* gene located on the short arm of chromosome 16. It is particularly frequent among Sephardic Jews, Armenians, Turks and Middle Eastern Arabs, where the prevalence can reach 1/2000 to 1/1000.

Recent publications described its frequent association with other diseases and/or syndromes, particularly those of autoimmune, genetic, and autoinflammatory origin.

The objective of this review is to familiarize healthcare professionals with the main associations to look for in patients followed for FMF. The early detection of these associations makes it possible to improve the management and the prognosis of patients with FMF.

Key words: familial Mediterranean fever, periodic fever, pyrin, chronic inflammation, *MEFV* gene.

Introduction

Familial Mediterranean fever (FMF) is an autoinflammatory disease with genetic transmission described in 1945 [1] and its characteristic mutation identified in 1992 [2, 3]. It is particularly frequent among Sephardic Jews, Armenians, Turks and Middle Eastern Arabs, where the prevalence can reach 1/2000 among Jews and 1/1000 for Turks and Armenians [1, 4].

It is caused by a point mutation in the *MEFV* gene located on the short arm of chromosome 16 [1, 4, 5]. This mutation leads to the production of a mutant protein pyrin and thus causes disruption of the normal inflammatory response and cellular apoptosis [4, 5].

Clinically, FMF is characterized by painful paroxysmal and febrile peritoneal, pleural, and joint attacks, which can be associated with various systemic manifestations: cutaneous, neurological, cardiac, urogenital and hematopoietic [1–4].

Several diagnostic criteria, the most used of which are those of Livneh [6], help in the positive diagnosis of this disease, but the confirmation is genetic, based on the demonstration of mutations in the *MEFV* gene, of which

there are more than twenty. The most frequent of these mutations are M694V, M694I, V726A, and E148Q [4, 7].

Recent publications have described its frequent association with other diseases and/or syndromes, the common denominators of which are genetic predisposition, immune dysfunction, and auto-inflammation. It is therefore important for health professionals to know these associations in order to detect them in time and initiate appropriate care without delay. This approach is the only guarantee to improve the prognosis of this disease, often reserved.

Familial Mediterranean fever and its most frequent associations

Several associations have been described with FMF; the most frequent are spondyloarthropathies, systemic vasculitis, and inflammatory bowel disease (IBD).

Familial Mediterranean fever and spondyloarthropathies

A significant association between FMF and spondyloarthropathies in general and ankylosing spondylitis in

Address for correspondence:

Salem Bouomrani, Sfax Faculty of Medicine, University of Sfax, Bd Majida Boulila, Sfax 3029, Tunisia, e-mail: salembouomrani@yahoo.fr

Submitted: 12.05.2020; Accepted: 8.06.2020

particular has been reported by several authors [8–11]. In fact, 7.5% of subjects with FMF in the Akar et al. [9] series had associated ankylosing spondylitis. The relative risk (RR) of having spondyloarthropathy and specifically ankylosing spondylitis was also very high in the first degree parents of a subject with FMF compared to the general population (PR = 3.3 and 2.9 respectively) [11].

This association is reinforced by the particular frequency of enthesopathies in subjects with FMF [12] suggesting that the mutant *MEFV* gene could partially promote such an association [11, 13].

In addition, a statistically significant association has been demonstrated between mutations in the *MEFV* gene, in particular that of type M694V, and ankylosing spondylitis (even without known FMF): 10.7% vs. only 4.2% in the general population, $p = 0.06$ in the series of Yigit et al. [10] and 22.3% vs. only 9.7% for all “nonsense” mutations of the *MEFV* gene in the Cosan et al. series [14].

Familial Mediterranean fever and systemic vasculitis

The most frequent association is with Henoch-Schönlein syndrome or purpura (HSS) [15, 16]. In fact, in the series of 68 patients with HSS of Altug et al. [17], 26% had an associated FMF. This particularly frequent association between HSS and FMF strongly suggests that *MEFV* could be the causative gene for HSS, the etiopathogenesis of which has not yet been elucidated [18]: in the series of 72 children with HSS of Dogan et al. [19], 11 (14.4%) had the *MEFV* gene mutation in its heterozygous form, 5 (6.6%) in its homozygous form, and 2 (2.6%) in its heterozygous composite form. In addition, seven of these patients presented at the same time two distinct mutations of the *MEFV* gene, which was clearly significant compared to the general population (9.2% vs. 1%) [19].

Other systemic vasculitides are also reported to be associated with FMF: panarteritis nodosa, protracted febrile myalgia syndrome (PFMS syndrome), Behçet's disease [11, 17, 20–22], and Takayasu arteritis [23, 24].

To approach the mechanism of these associations, the study by Aytakin et al. [22] seems very interesting by revealing a particularly high frequency of vascular anomalies on capillaroscopy during FMF (capillary enlargement and microhemorrhages). These anomalies which characterize systemic vasculitides suggest systemic angiitis of the small vessels as a mediator of FMF clinical symptoms [22]. This hypothesis is reinforced by the original observation of Girisgen et al. [16] reporting the concomitant association of FMF with two systemic vasculitides (HSS and panarteritis nodosa).

In summary, among the systemic vasculitides associated with FMF, HSS and classic panarteritis nodo-

sa are the most frequent [11, 24]; next come the PFMS syndrome and Behçet's disease [11]. The other systemic vasculitides are more exceptional [23, 24].

Familial Mediterranean fever and inflammatory bowel disease

Inflammatory bowel disease (Crohn's disease and ulcerative colitis) are far from rare during FMF [11, 25–28]. In fact, 22 (15.4%) in the series of 78 patients suffering from FMF of Beser OF et al. [26] had concomitant endoscopically and histologically confirmed IBD. Likewise, the presence of the *MEFV* gene mutation was significantly more often found in patients with histologically confirmed IBD: 28% in ulcerative colitis, and 22.6% in Crohn's disease vs. only 9.9% in the healthy controls; $p = 0.006$ [27].

Familial Mediterranean fever and multiple sclerosis

This association reported by several authors [11, 29–32] but too long controversial [33] has just been recently proven by the large national study of Yahalom et al. (one of the largest series of FMF) where the frequency of multiple sclerosis in patients with FMF was 0.075% = three times that of the general population ($p = 0.0057$); in addition, the presence of the M694V mutation worsened the clinical picture of multiple sclerosis in these patients [34].

On the other hand, the study of Unal et al. [29] had shown a significant frequency of the mutation of the *MEFV* gene in patients with multiple sclerosis compared to the general population: 38% vs. only 11%, $p < 0.0001$.

These findings suggest the mutation of the *MEFV* gene as a potential predisposing factor for multiple sclerosis, in particular the E148Q type mutation [32].

Familial Mediterranean fever and rheumatoid arthritis

This association is increasingly reported in the world literature [35–39], and a significantly higher frequency of anti-citrullinated protein antibodies (ACPA) (specific for rheumatoid arthritis) has been found in patients with FMF: 14.5% vs. only 4.7% in the general population (3 times higher); their presence was significantly associated with arthritis during FMF [40].

Familial Mediterranean fever and juvenile idiopathic arthritis

This association is also far from rare. It has been reported by several authors [41–43]. It has been shown that the frequency of the specific M694V mutation of the *MEFV* gene during JIA is around 10%, and its pres-

Table I. Main clinical associations of familial Mediterranean fever presented in the literature

Most frequent associations	Rarer associations
Spondyloarthropathies, specifically ankylosing spondylitis	Systemic lupus erythematosus
Systemic vasculitis: Henoch-Schönlein syndrome, panarteritis nodosa, protracted febrile myalgia syndrome (PFMS syndrome), Behçet's disease, Takayasu arteritis	Sarcoidosis and Blau syndrome
Inflammatory bowel disease: Crohn's disease and ulcerative colitis	Kartagener syndrome
Multiple sclerosis	Gitelman syndrome
Rheumatoid arthritis and juvenile idiopathic arthritis	Fabry syndrome
Non-alcoholic fatty liver disease	Celiac disease
	Polymyositis and juvenile dermatomyositis
	Beta-thalassemia
	Atrophic polychondritis
	Fibromyalgia
	Eosinophilic gastroenteritis
	Tumor necrosis factor receptor-1 syndrome (TRAPS syndrome)

ence makes juvenile idiopathic arthritis (JIA) more severe and more resistant to the usual treatment requiring the use of biotherapy [41]. Also the presence of all types of *MEFV* mutations was significantly higher in JIA compared to the general population: 14.28% vs. only 5%, $p < 0.01\%$ [41].

Therefore, it is recommended to screen for *MEFV* gene mutations in children presumed to have JIA, especially those of male gender, in the case of enthesopathies and negative anti-nuclear antibodies [44].

Familial Mediterranean fever and non-alcoholic fatty liver disease

In the series of Rimar et al. [45], 74% of patients with FMF presented non-alcoholic fatty liver disease (NAFLD) ranging from simple hepatic steatosis (5/21 patients) to non-alcoholic steatohepatitis (NASH): 3/21 patients, and cirrhosis on NASH: 7/21 patients, and this apart from any underlying metabolic syndrome, suggesting a strong association between FMF and NAFLD.

Familial Mediterranean fever and its rarer associations

Other rarer associations have been reported, with: systemic lupus erythematosus [46-48], sarcoidosis [38], Blau syndrome [49], Kartagener syndrome [50], Gitelman syndrome [51], Fabry syndrome [52], celiac disease [53], polymyositis [54], juvenile dermatomyositis [55], beta-thalassemia [56], atrophic polychondritis [57], fibromyalgia [58], eosinophilic gastroenteritis [59], and tumor necrosis factor receptor-1 syndrome (TRAPS syndrome) [60].

Table I summarizes the main associations found in the literature with FMF.

Conclusions

Knowledge of these associations, especially the most common, is very useful for healthcare professionals caring for patients with FMF. Their diagnosis is not always easy since many of these diseases can cause the same symptoms or have similar clinical presentations.

The early detection of these associations makes it possible to improve the management and the prognosis of patients with FMF.

The authors declare no conflict of interest.

References

1. Etem E, Deveci SD, Erol D, et al. Familial Mediterranean Fever: a retrospective clinical and molecular study in the East of Anatolia region of Turkey. *Open Rheum J* 2010; 4: 1-6, DOI: 10.2174/1874312901004010001.
2. The French FMF consortium. A candidate gene for familial Mediterranean fever. *Nat Genet* 1997; 17: 25-31.
3. International FMF Consortium: Ancient missense mutations in a new member of the RoRet gene family are likely to cause familial Mediterranean fever. *Cell* 1997; 90: 797-807, DOI: 10.1016/s0092-8674(00)80539-5.
4. Medlej-Hashim M, Loiselet J, Lefranc G, Mégarbané A. Familial Mediterranean Fever (FMF): from diagnosis to treatment. *Sante* 2004; 14: 261-266.
5. Tidow N, Chen X, Müller C, et al. Hematopoietic-specific expression of *MEFV*, the gene mutated in familial Mediterranean fever, and subcellular localization of its corresponding protein, pyrin. *Blood* 2000; 95: 1451-1455.

6. Livneh A, Laguevitz P, Zemer D, et al. Criteria for the diagnosis of familial Mediterranean fever. *Arthritis Rheum* 1997; 40: 1879-1885, DOI: 10.1002/art.1780401023.
7. Touitou I. The spectrum of Familial Mediterranean Fever (FMF) mutations. *Eur J Hum Genet* 2001; 9: 473-483, DOI: 10.1038/sj.ejhg.5200658.
8. Estublier C, Stankovic Stojanovic K, Bergerot JF, et al. Myositis in a patient with familial Mediterranean fever and spondyloarthritis successfully treated with anakinra. *Joint Bone Spine* 2013; 80: 645-649, DOI: 10.1016/j.jbspin.2013.03.004.
9. Akar S, Soysal O, Balci A, et al. High prevalence of spondyloarthritis and ankylosing spondylitis among familial Mediterranean fever patients and their first-degree relatives: further evidence for the connection. *Arthritis Res Ther* 2013; 15: R21, DOI: 10.1186/ar4154.
10. Yigit S, Inanir A, Karakus N, et al. Common Mediterranean fever (MEFV) gene mutations associated with ankylosing spondylitis in Turkish population. *Dis Markers* 2012; 33: 113-118, DOI: 10.1155/2012/890214.
11. Aksu K, Keser G. Coexistence of vasculitides with familial Mediterranean fever. *Rheumatol Int* 2011; 31: 1263-1274, DOI: 10.1007/s00296-011-1840-z.
12. Ozkan F, Cetin GY, Inci MF, et al. Increased enthesopathy in patients with familial Mediterranean fever: evaluation with a new sonographic enthesitis index. *J Ultrasound Med* 2013; 32: 325-332, DOI: 10.7863/jum.2013.32.2.325.
13. Tufan A, Mercan R, Tezcan ME, et al. Enthesopathy in patients with familial Mediterranean fever: increased prevalence in M694 V variant. *Rheumatol Int* 2013; 33: 1933-1937, DOI: 10.1007/s00296-013-2669-4.
14. Cosan F, Ustek D, Oku B, et al. Association of familial Mediterranean fever-related MEFV variations with ankylosing spondylitis. *Arthritis Rheum* 2010; 62: 3232-3236, DOI: 10.1002/art.27683.
15. Cabral M, Conde M, Brito MJ, et al. Protracted Febrile Myalgia Syndrome with Henoch-Schönlein Purpura: an atypical presentation of Familial Mediterranean Fever. *Acta Reumatol Port* 2011;36: 69-74.
16. Girisgen I, Sonmez F, Koseoglu K, et al. Polyarteritis nodosa and Henoch-Schönlein purpura nephritis in a child with familial Mediterranean fever: a case report. *Rheumatol Int* 2012; 32: 529-533, DOI: 10.1007/s00296-010-1766-x.
17. Altug U, Ensari C, Sayin DB, Ensari A. MEFV gene mutations in Henoch-Schönlein purpura. *Int J Rheum Dis* 2013; 16: 347-351, DOI: 10.1111/1756-185X.12072.
18. He X, Yu C, Zhao P, et al. The genetics of Henoch-Schönlein purpura: a systematic review and meta-analysis. *Rheumatol Int* 2013; 33: 1387-1395, DOI: 10.1007/s00296-012-2661-4.
19. Dogan CS, Akman S, Koyun M, et al. Prevalence and significance of the MEFV gene mutations in childhood Henoch-Schönlein purpura without FMF symptoms. *Rheumatol Int* 2013; 33: 377-380, DOI: 10.1007/s00296-012-2400-x.
20. Luger S, Harter PN, Mittelbronn M, et al. Brain stem infarction associated with familial Mediterranean fever and central nervous system vasculitis. *Clin Exp Rheumatol* 2013;31 (3 Suppl 77): 93-95.
21. Özçakar ZB, Yalçınkaya F. Vascular comorbidities in familial Mediterranean fever. *Rheumatol Int* 2011; 3: 1275-1281, DOI: 10.1007/s00296-011-1845-7.
22. AYTEKİN S, AYDIN F, AKPOLAT T, et al. Evaluation of nailfold capillaries in familial Mediterranean fever patients. *Clin Rheumatol* 2011; 30: 255-257, DOI: 10.1007/s10067-010-1609-x.
23. Alibaz-Oner F, Yılmaz N, Can M, Direskeneli H. A case of Takayasu's arteritis associated with familial Mediterranean fever. *Clin Exp Rheumatol* 2012; 30 (3 Suppl 72): S117.
24. Zihni FY, Kalfa M, Ocakçı PT, et al. Coexistence of Takayasu's arteritis with familial Mediterranean fever. *Rheumatol Int* 2012; 32: 1675-1678, DOI: 10.1007/s00296-011-1853-7.
25. Kuloğlu Z, Kansu A, Ustündağ G, et al. An infant with severe refractory Crohn's disease and homozygous MEFV mutation who dramatically responded to colchicine. *Rheumatol Int* 2012; 32: 783-785, DOI: 10.1007/s00296-009-1326-4.
26. Beşer OF, Kasapçopur O, Cokuğraş FC, et al. Association of inflammatory bowel disease with familial Mediterranean fever in Turkish children. *J Pediatr Gastroenterol Nutr* 2013; 56: 498-502, DOI: 10.1097/MPG.0b013e31827dd763.
27. Akyuz F, Besisik F, Ustek D, et al. Association of the MEFV gene variations with inflammatory bowel disease in Turkey. *J Clin Gastroenterol* 2013; 47: e23-27, DOI: 10.1097/MCG.0b013e3182597992.
28. Uslu N, Demir H, Balta G, et al. Hemophagocytic syndrome in a child with severe Crohn's disease and familial Mediterranean fever. *J Crohns Colitis* 2010; 4: 341-344, DOI: 10.1016/j.crohns.2009.12.005.
29. Unal A, Dursun A, Emre U, et al. Evaluation of common mutations in the Mediterranean fever gene in Multiple Sclerosis patients: Is it a susceptibility gene? *J Neurol Sci* 2010; 294: 38-42, DOI: 10.1016/j.jns.2010.04.008.
30. Unal A, Emre U, Dursun A, Aydemir S. The co-incidence of multiple sclerosis in a patient with familial Mediterranean fever. *Neurol India* 2009; 57: 672-673, DOI: 10.4103/0028-3886.57790.
31. Alpaycı M, Bozan N, Erdem S, et al. The possible underlying pathophysiological mechanisms for development of multiple sclerosis in familial Mediterranean fever. *Med Hypotheses* 2012; 78: 717-720, DOI: 10.1016/j.mehy.2012.02.017.
32. Kümpfel T, Gerdes LA, Wacker T, et al. Familial Mediterranean fever-associated mutation pyrin E148Q as a potential risk factor for multiple sclerosis. *Mult Scler* 2012; 18: 1229-1238, DOI: 10.1177/1352458512437813.
33. Yahalom G, Kivity S, Lidar M, et al. Familial Mediterranean fever (FMF) and multiple sclerosis: an association study in one of the world's largest FMF cohorts. *Eur J Neurol* 2011; 18: 1146-1150, DOI: 10.1111/j.1468-1331.2011.03356.x.
34. Bouomrani S, Guerhazi M. Multiple Sclerosis and Familial Mediterranean Fever: Myth or Reality? *Ann Mult Scler Res* 2019; 1: 1003.
35. Mori S, Yonemura K, Migita K. Familial Mediterranean fever occurring in an elderly Japanese woman with recent-onset rheumatoid arthritis. *Intern Med* 2013; 52: 385-388, DOI: 10.2169/internalmedicine.52.9102.
36. Koca SS, Etem EO, Isik B, et al. Prevalence and significance of MEFV gene mutations in a cohort of patients with rheumatoid

- arthritis. *Joint Bone Spine* 2010; 77: 32-35, DOI: 10.1016/j.jbspin.2009.08.006.
37. Migita K, Abiru S, Sasaki O, et al. Coexistence of familial Mediterranean fever and rheumatoid arthritis. *Mod Rheumatol* 2014; 24: 212-216, DOI: 10.3109/14397595.2013.852843.
38. Erten S, Erzurum C, Kosker TA, et al. Two cases of familial Mediterranean fever associated with sarcoidosis (Lofgren's syndrome) and rheumatoid arthritis. *Int J Rheum Dis* 2013; 16: 373-375, DOI: 10.1111/1756-185X.12004.
39. Migita K, Abiru S, Sasaki O, et al. Coexistence of familial Mediterranean fever and rheumatoid arthritis. *Mod Rheumatol* 2014; 24: 212-216, DOI: 10.3109/14397595.2013.852843.
40. Ceri M, Unverdi S, Altay M, et al. Anti-cyclic citrullinated peptides positivity rate in patients with familial Mediterranean fever. *Clin Exp Rheumatol* 2010; 28 (4 Suppl 60): S58-61.
41. Ayaz NA, Ozen S, Bilginer Y, et al. MEFV mutations in systemic onset juvenile idiopathic arthritis. *Rheumatology (Oxford)* 2009; 48: 23-25, DOI: 10.1093/rheumatology/ken409.
42. Kaya S, Kaptanoglu E, Elden H, Hizmetli S. Coexistence of familial Mediterranean fever and juvenile idiopathic arthritis with osteoporosis successfully treated with etanercept. *Intern Med* 2010; 49: 619-622, DOI: 10.2169/internalmedicine.49.2777 .
43. Cantarini L, Lucherini OM, Simonini G, et al. Systemic-onset juvenile idiopathic arthritis complicated by early onset amyloidosis in a patient carrying a mutation in the MEFV gene. *Rheumatol Int* 2012; 32: 465-467, DOI: 10.1007/s00296-009-1331-7.
44. Comak E, Dogan CS, Akman S, et al. MEFV gene mutations in Turkish children with juvenile idiopathic arthritis. *Eur J Pediatr* 2013; 172: 1061-1067, DOI: 10.1007/s00431-013-2003-x.
45. Rimar D, Rosner I, Rozenbaum M, Zuckerman E. Familial Mediterranean fever: an association with non-alcoholic fatty liver disease. *Clin Rheumatol* 2011; 30: 987-991, DOI: 10.1007/s10067-011-1718-1.
46. Inamo Y, Hayashi R, Saito K, et al. A Japanese pediatric patient with coexisting systemic lupus erythematosus and familial Mediterranean fever. *Lupus* 2013; 22: 1056-1059, DOI: 10.1177/0961203313497117.
47. Matsuda M, Kishida D, Tsuchiya-Suzuki A, et al. Periodic peritonitis due to familial Mediterranean fever in a patient with systemic lupus erythematosus. *Intern Med* 2010; 49: 2259-2262, DOI: 10.2169/internalmedicine.49.4043.
48. Yildiz G, Kayatas M, Uygun Y, et al. Coexistence of systemic lupus erythematosus and familial Mediterranean fever. *Intern Med* 2010; 49: 767-769, DOI: 10.2169/internalmedicine.49.3102.
49. Oğuz ID, Gönül M, Çakmak SK, et al. A case of Beau's lines associated with familial Mediterranean fever. *Int J Dermatol* 2014; 53: e121-2, DOI: 10.1111/j.1365-4632.2012.05618.x.
50. Cetin D, Genç Çetin B, Sentürk T, et al. Coexistence of two rare genetic disorders: Kartagener syndrome and familial Mediterranean fever. *Mod Rheumatol* 2015; 25: 312-314, DOI: 10.3109/14397595.2013.874756.
51. Erten S, Ceylan GG, Altunoğlu A. Concomitance of Gitelman syndrome and familial Mediterranean fever: A rare case presentation. *Ren Fail* 2012; 34: 1333-1334, DOI: 10.3109/0886022X.2012.718950.
52. Zizzo C, Colomba P, Albegiani G, et al. Misdiagnosis of familial Mediterranean fever in patients with Anderson-Fabry disease. *Clin Genet* 2013; 83: 576-581, DOI: 10.1111/j.1399-0004.2012.01940.x.
53. Krums LM, Golovanova EV, Khomeriki SG, et al. Celiac disease is a 33-year-old man with periodic disease. *Eksp Klin Gastroenterol* 2012; 3: 114-117.
54. Eguchi M, Miyashita T, Shirouzu H, et al. Coexistence of polymyositis and familial Mediterranean fever. *Mod Rheumatol* 2013; 23: 374-378, DOI: 10.1007/s10165-012-0649-2.
55. Bouomrani S, Mesfar R, Ben Ayed M, et al. Juvenile dermatomyositis associated to Familial Mediterranean Fever. *Am J Med Case Rep* 2019; 7:170-172, DOI: 10.12691/ajmer-7-8-5.
56. Aslan D. Familial Mediterranean fever with a single MEFV mutation: can a deletion resulting in α -thalassemia be the cause? *J Hum Genet* 2011; 56: 169-171, DOI: 10.1038/jhg.2010.160.
57. Miller EB, Friedman JA, Lahav Y, Landau Z. Relapsing polychondritis and familial Mediterranean fever – an association. *Clin Rheumatol* 2011; 30: 711-713, DOI: 10.1007/s10067-010-1673-2.
58. Feng J, Zhang Z, Li W, et al. Missense mutations in the MEFV gene are associated with fibromyalgia syndrome and correlate with elevated IL-1beta plasma levels. *PLoS One* 2009; 4: e8480, DOI: 10.1371/journal.pone.0008480.
59. Kocak G, Kocak E, Yilmaz SR, et al. MEFV gene mutations in a patient with eosinophilic gastroenteritis. *South Med J* 2010; 103: 1170-1172, DOI: 10.1097/SMJ.0b013e3181f468b4.
60. Clementi A, Cruz ND, Granata A, et al. Secondary amyloidosis in a patient carrying mutations in the familial Mediterranean fever (FMF) and tumour necrosis factor receptor-1 syndrome (TRAPS) genes. *Clin Kidney J* 2013; 6: 613-617, DOI: 10.1093/ckj/sft112.w